



	Ref	Description	Dimensions	Packaging
	0511	Flexible Surgical Dressing	5.3" x 11" (13cm x 28cm)	5 Per Box 2 Boxes Per Case
	1511	Flexible Silver Surgical Dressing	5.3" x 11" (13cm x 28cm)	5 Per Box 2 Boxes Per Case
	0512	Surgical Dressing	5.5" x 12.75" (14cm x 32cm)	5 Per Box 2 Boxes Per Case
	1512	Silver Surgical Dressing	5.5" x 12.75" (14cm x 32cm)	5 Per Box 2 Boxes Per Case

PolyMem Surgical dressings are latex-free.

INDICATIONS FOR USE

For the management of partial and full thickness surgical wounds healing by primary intention.

Available through most major medical supply distributors.
This document is meant for general informational purposes only. See individual product literature for specific indication and instructions for use.
We strongly recommend that you consult your payer organization with regard to its reimbursement policies.

References:

1. Beitz AJ, Newman A, Kahn AR, Ruggles T, Eikmeier L. A Polymeric Membrane Dressing with Antinociceptive Properties: Analysis with a Rodent Model of Stab Wound Secondary Hyperalgesia. The Journal of Pain. Feb 2004;5(1):38-47
2. Knight KL. Chapter 3. Inflammation and Wound Repair. In Cryotherapy in Sport Injury Management. Human Kinetics. 1995. Champaign, IL
3. Merrick MA. Secondary injury after musculoskeletal trauma: a review and update. Journal of Athletic Training 2002;37(2):209-217
4. Clay CS, Chen WYJ. Wound pain: the need for a more understanding approach. Journal of Wound Care. April 2005;14(4):181-184
5. Abraham SE. Pain Management in wound care. Podiatry Management. June/July 2006;165-168
6. Wulf H, Baron R. The Theory of Pain in European Wound Management Association Position Document Pain at Wound Dressing Changes, Medical Education Partnership, London UK, 2002; page 8-11
7. Levine JD, Reichling DB. Chapter 2 Peripheral Mechanisms of Inflammatory Pain. In Wall PD, Melzak R, Editors. Textbook of Pain. 4th edition. Edinburgh, UK: Churchill Livingstone, 1999. pages 59-84.
8. Fields HL. Chapter 1 Introduction & Chapter 2 The Peripheral Pain Sensory System In Pain New York: McGraw-Hill 1987 pages 1-40
9. Brook I. Clinical clues to diagnosis of anaerobic infections <http://www.medscape.com/viewarticle/495997> accessed July 18, 2010
10. Wolcott RD, Rhoads DD, Dowd SE. Biofilms and chronic wound inflammation. Journal of Wound Care 2008; 17(8):333-341
11. Dow G, Browne A, Sibbald R.G. Infection in chronic wounds: Controversies in diagnosis and treatment. Ostomy/Wound Management 1999;45(8):23-40
12. Dawson N, Lewis C. Total joint replacement surgical site infections eliminated by using multifunctional dressing. 900 cases report over 4 years. Poster. Australian College of operating room Nurses (ACORN) May 19-22, 2010. Perth, Australia
13. Rosevear C, Dott J, Lazarus R. Reducing risk of post-operative complications after joint replacement surgery. Poster. Australian College of operating room Nurses (ACORN) May 19-22, 2010. Perth, Australia
14. Schmid P. Reduction of edema and pain on surgical wounds with polymeric membrane dressings. Poster. European Wound Management Association (EWMA) Poster #P317. May 26-28, 2010. Geneva, Switzerland
15. Burkhard R. Efficient reduction of swelling and bruising on severe sports injuries when using polymeric membrane dressings. Poster. European Wound Management Association (EWMA) Poster #P218. May 26-28, 2010. Geneva, Switzerland
16. Burd A, Kwok CH, Hung SC, Chan HS, Gu H, Lam WK, Huang L. A Comparative Study of the Cytotoxicity of Silver-Based Dressings in Monolayer Cell, Tissue Explant, and Animal Models. Wound Repair and Regeneration. Jan-Feb 2007; 15(1):94-104

This document is meant for general informational purposes only. See individual product literature for specific indication and instructions for use.

PolyMem, PolyMem Silver, PolyMem Surgical, PolyMem Silver Surgical, More Healing • Less Pain, interlocking circles design, Ferris and FMC Ferris and design are marks owned by or licensed to Ferris. The marks may be registered or pending in the US Patent and Trademark Office and in other countries. Other marks are the property of their respective owners.
© 2013 Ferris Mfg. Corp. MKL-627, REV.0, 0413

♻️ 30% post-consumer material



Ferris Mfg. Corp.

5133 Northeast Parkway, Fort Worth, TX 76106 U.S.A.

Toll Free U.S.A.: 800.POLYMEM (765.9636)

International: +1 817-900-1301

www.polymem.com

Reduce Inflammation, Increase Healing

Focus inflammatory response for maximum benefit, speed healing, and improve outcomes with a revolutionary dressing for surgical incision sites




POLYMEM®—THE PROVEN INTELLIGENT DRESSING

PolyMem dressings have helped clinicians successfully manage millions of wounds worldwide. The dressings protect the wound and ensure moist wound healing by continuously cleansing, filling, absorbing and moistening wounds, regardless if they are healing by primary or secondary intention.

Activated by wound fluid...

- The PolyMem dressing will expand and conform to the wound.
- The mild, non-ionic, non-toxic, tissue-friendly wound cleanser/surfactant and the glycerin incorporated in the dressing will be released to the wound bed/incision site, while the starch co-polymer and the foam will bind fluid in the dressing.
- The semi-permeable film cover helps maintain appropriate moisture balance.



PolyMem Surgical has the ability to rapidly reduce inflammation, resulting in the following benefits:

CLINICIANS


- Improved compliance post-operatively
- Less chance of infection and wound breakdown
- Less reliance on analgesia
- Overall cost-benefit

PATIENTS


- Reduced pain
- Reduced edema and swelling
- Improved rehabilitation
- Overall improved clinical outcome




PolyMem helps reduce spread of inflammatory reaction into surrounding, uninjured areas



(A) Incision only



(B) Incision with gauze



(C) Incision with PolyMem

This series of images shows the width of the spread of the inflammatory cells, in muscle, around an incision. The dark portion of the scale in each image (each segment is 100µm) represents the spread of the zone of the inflammatory reaction around the center line of the incision. In images A and B, there is no difference in the spread of the inflammatory reaction around the center of the injury. In image C, notice how PolyMem reduces the spread of the inflammation into the surrounding tissue. Statistically, PolyMem reduces the spread of the inflammatory reaction into the surrounding undamaged tissue by approximately 25 percent.¹

Focused inflammation improves outcomes

PolyMem Surgical dressings are specifically designed to focus inflammation to where it is needed—in the operated tissues.¹

Focusing the inflammatory process helps reduce secondary cell damage and pain caused by the typical swelling and bruising usually observed beyond the operative site.^{1,2,3}

PolyMem has been shown to reduce secondary cell damage by reducing the recruitment of adjacent inflammatory nerve endings (also referred to as nociceptors or free nerve endings).¹ These populous nerve endings, found in the epidermis, dermis, muscle, joints and viscera, are responsible for triggering and spreading the inflammatory reaction into surrounding uninjured tissues.^{4,5,6,7,8}

The spreading of inflammation is often clinically evidenced by increased temperature, bruising, swelling, increased sensitivity to stimuli, and pain beyond the immediate zone of injury.^{5,7}

PolyMem, when placed on open or closed injuries, is believed to provide its benefits by changing the nerve responses both locally, at the injury site, and centrally at the level of the dorsal root ganglion and spinal cord.¹

Minimize infection and blistering

By helping to reduce swelling and bruising, PolyMem improves blood flow to the surgical site and removal of cellular debris. All of these actions help to reduce the immediate and long-term risk of infection.^{2,3,9,10,11} PolyMem has been shown to help reduce the risk of surgical site and joint infection and virtually eliminate blistering, while dramatically reducing patients' swelling and pain.^{12,13,14,15}

No infections in 900 consecutive total joint replacements.¹⁸

These dressings, applied in the operative theater, are designed to be left in place for three to five days so the incision is not exposed to potential contamination during a dressing change or visualization prior to epithelialization completion.¹³

For additional antibacterial action, PolyMem Silver Surgical dressings can be used to manage incision sites.* The dressings contain high-purity, nanocrystalline silver particles which are universally distributed throughout and bound into the dressing membrane. The dressings have been found to be one of the best for use on healing tissues.¹⁶ Other commonly specified silver dressings have been found to damage the key cells required for healing.¹⁶

* In tests for antimicrobial effectiveness using several in-vitro methods, PolyMem Silver dressings killed at least 99.9% of the entire population of each organism tested. The bacteria and fungi tested included Klebsiella pneumoniae (ATCC# 4352), Pseudomonas aeruginosa (ATCC# 9027), Enterococcus faecalis (VRE) (ATCC# 51575), Candida albicans (ATCC# 10231), Staphylococcus aureus (MRSA) (ATCC# 33591) and Staphylococcus aureus (ATCC#6538). The organisms chosen demonstrate the antimicrobial actions of the silver formulation on relevant, representative organisms found in clinical settings.